

INTRODUCTION TO THE TORIC MORI THEORY

OSAMU FUJINO AND HIROSHI SATO

ABSTRACT. The main purpose of this paper is to give a simple and non-combinatorial proof of the toric Mori theory. Here, the toric Mori theory means the (log) Minimal Model Program (MMP, for short) for toric varieties. We minimize the arguments on fans and their decompositions. We recommend this paper to the following people:

- (A) those who are uncomfortable with manipulating fans and their decompositions,
- (B) those who are familiar with toric geometry but not with the MMP.

People in the category (A) will be relieved from tedious combinatorial arguments in several problems. Those in the category (B) will discover the potential of the toric Mori theory.

As applications, we treat the Zariski decomposition on toric varieties and the partial resolutions of non-degenerate hypersurface singularities. By these applications, the reader will learn to use the toric Mori theory.

CONTENTS

1. Introduction	2
2. Preliminaries	3
2.1. Toric varieties	3
2.2. Singularities of pairs	6
3. Toric Mori theory	7
4. Proof of the toric Mori theory	8
5. On the Zariski decomposition	14
6. Application to hypersurface singularities	16
References	19

1991 *Mathematics Subject Classification.* Primary 14M25; Secondary 14E30, 14E15.

Key words and phrases. Toric varieties, Mori theory, Minimal Model Program, Zariski decomposition, hypersurface singularities.

The second author is partly supported by the Grant-in-Aid for JSPS Fellows, The Ministry of Education, Science, Sports and Culture, Japan.

1. INTRODUCTION

The main purpose of this paper is to give a simple and non-combinatorial proof of the toric Mori theory. Here, the toric Mori theory means the (log) Minimal Model Program (MMP, for short) for toric varieties.

In his famous and beautiful paper [R], Reid carried out the toric Mori theory under the assumption that the variety is *complete*. His arguments are combinatorial. Thus, it is not so obvious whether we can remove the completeness assumption from his paper. We quote his idea from [R].

(0.3) *Remarks.* The hypothesis that A is complete is not essential; it can be reduced to the projective case, or possibly eliminated by a careful (and rather tedious) rephrasing of the arguments of §§1–3. The projectivity hypothesis on f is needed in order for the statement of (0.2) to make sense, since without projectivity the cone $\text{NE}(V/A)$ will usually not have any extremal rays.

We prefer not to simply rephrase his approach, which entails tedious combinatorial arguments. Instead, our proof, which is independent of Reid's proof, heavily relies on the general machinery of the Minimal Model Program and the special properties of toric varieties. Thus, our proof works without the completeness assumption.

For the details of the toric Mori theory, see [O, §2.5], [KMM, §5.2], [OP], [L], [I2], [Ma, Chapter 14], [W], and [Fj1]. Matsuki [Ma] corrected some minor errors in [R] and pointed out some ambiguities in [R] and [KMM]. See Remarks 14-1-3 (ii), 14-2-3, and 14-2-7 in [Ma]. We believe that these remarks help the reader to understand [R]. We recommend that the reader compare this paper with [Ma, Chapter 14]. Shokurov [Sh] treats the MMP for toric varieties in a non-combinatorial way (see [Sh, Example 3]). His arguments are quite different from ours. For the more advanced topics of the toric Mori theory, see [Fj2].

For the outline of the general MMP, which is still conjectural in dimension ≥ 4 , see [KMM, Introduction] or [KM, 2.14, §3.7].

We note that the Zariski decomposition on toric varieties has already been treated by various researchers. The reason we treat it here is to show that the Zariski decomposition on toric varieties is an easy consequence of the toric Mori theory. The partial resolutions of non-degenerate hypersurface singularities were treated by Ishii, who divided the cone by the data of the Newton polytope. One of her proofs contains a gap (see Remark 6.6), but the results themselves are correct. Her results also become easy consequences of the toric Mori theory.

Note that we cannot recover combinatorial aspects of [R] and [I1] by our method. So, this paper does not depreciate [R] and [I1].

Almost all the results in this paper are more or less known to the experts. However, some of them were not stated explicitly before. Some of the proofs that we give in this paper are new and much simpler than the known ones. We hope that this paper will help the reader to understand the toric Mori theory.

We summarize the contents of this paper: In Section 2, we fix the notation and collect basic results. Section 3 explains the toric Mori theory. Section 4 is the main part of this paper. There, we give a simple and non-combinatorial proof of the toric Mori theory. In Section 5, we consider the Zariski decomposition on toric varieties. In Section 6, we apply the toric Mori theory to the study of the partial resolutions of non-degenerate hypersurface singularities. We reprove Ishii's results.

Notation. Here is a list of some of the standard notation we use.

- (1) For a real number d , its *round down* is the largest integer $\leq d$. It is denoted by $\lfloor d \rfloor$. If $D = \sum d_i D_i$ is a divisor with real coefficients and the D_i are distinct prime divisors, then we define the *round down* of D as $\lfloor D \rfloor := \sum \lfloor d_i \rfloor D_i$.
- (2) Let $f : X \rightarrow Y$ be a proper birational morphism between normal varieties. Then, f is said to be *small* if f is an isomorphism in codimension one.
- (3) The symbol $\mathbb{Z}_{\geq 0}$ (resp. $\mathbb{Q}_{\geq 0}$, $\mathbb{R}_{\geq 0}$) denotes the set of non-negative integers (resp. rational numbers, real numbers).

We will work over an algebraically closed field k throughout this paper. The characteristic of k is arbitrary from Section 2 to Section 5 unless otherwise stated. In Section 6, we assume that k is the complex number field \mathbb{C} .

Acknowledgments. We would like to thank Professors Shigefumi Mori and Tatsuhiro Minagawa for fruitful discussions and useful comments. We are grateful to Professor Shihoko Ishii, who kindly answered our questions about the proof of [I1, Theorem 3.1] and gave us several useful comments. Finally, we thank Professor Tadao Oda, whose comments on the preliminary version made this paper more readable.

2. PRELIMINARIES

2.1. Toric varieties. In this subsection, we recall the basic notion of toric varieties and fix the notation. For the basic results about toric varieties, see [KKMS], [D], [O], and [Fl].

2.1. Let $N \simeq \mathbb{Z}^n$ be a lattice of rank n . A toric variety $X(\Delta)$ is associated to a fan Δ , a finite collection of convex cones $\sigma \in \Delta$ in $N_{\mathbb{R}} := N \otimes_{\mathbb{Z}} \mathbb{R}$ satisfying the following:

- (i) Each convex cone $\sigma \in \Delta$ is rational polyhedral in the sense that there are finitely many $v_1, \dots, v_s \in N \subset N_{\mathbb{R}}$ such that

$$\sigma = \{r_1 v_1 + \dots + r_s v_s; r_i \in \mathbb{R}_{\geq 0} \text{ for all } i\}$$

and is strongly convex in the sense that

$$\sigma \cap (-\sigma) = \{0\}.$$

- (ii) Each face τ of a convex cone $\sigma \in \Delta$ again belongs to Δ .
- (iii) The intersection of two cones in Δ is a face of each.

Definition 2.2. The *dimension* $\dim \sigma$ of σ is the dimension of the linear space $\mathbb{R} \cdot \sigma = \sigma + (-\sigma)$ spanned by σ .

We define the sublattice N_{σ} of N generated (as a subgroup) by $\sigma \cap N$ as follows:

$$N_{\sigma} := \sigma \cap N + (-\sigma \cap N).$$

The *star* of a cone τ can be defined abstractly as the set of cones σ in Δ that contain τ as a face. For any such cone σ , the image

$$\bar{\sigma} = (\sigma + (N_{\tau})_{\mathbb{R}}) / (N_{\tau})_{\mathbb{R}} \subset N(\tau)_{\mathbb{R}}$$

by the projection $N \rightarrow N(\tau) := N / N_{\tau}$ is a cone in $N(\tau)$. These cones $\{\bar{\sigma}; \tau \prec \sigma\}$ form a fan in $N(\tau)$, and we denote this fan by $\text{Star}(\tau)$. We set $V(\tau) = X(\text{Star}(\tau))$. It is well-known that $V(\tau)$ is an $(n - k)$ -dimensional closed toric subvariety of $X(\Delta)$, where $\dim \tau = k$. If $\dim V(\tau) = 1$ (resp. $n - 1$), then we call $V(\tau)$ a *torus invariant curve* (resp. *torus invariant divisor*). For details on the correspondence between τ and $V(\tau)$, see, for instance, [Fl, 3.1 Orbits].

Definition 2.3 (\mathbb{Q} -Cartier divisor and \mathbb{Q} -factoriality). Let $D = \sum d_i D_i$ be a \mathbb{Q} -divisor on a normal variety X , that is, $d_i \in \mathbb{Q}$ and D_i is a prime divisor on X for every i . Then D is \mathbb{Q} -Cartier if there exists a positive integer m such that mD is a Cartier divisor. A normal variety X is said to be \mathbb{Q} -factorial if every prime divisor D on X is \mathbb{Q} -Cartier.

The next lemma is well-known. See, for example, [R, (1.9)] or [Ma, Lemma 14-1-1].

Lemma 2.4. *A toric variety $X(\Delta)$ is \mathbb{Q} -factorial if and only if each cone $\sigma \in \Delta$ is simplicial.*

The following remarks are easy but important.

Remark 2.5. Let D be a Cartier (resp. \mathbb{Q} -Cartier) divisor on a toric variety. Then D is linearly (resp. \mathbb{Q} -linearly) equivalent to a torus invariant divisor (resp. \mathbb{Q} -divisor).

Remark 2.6 (cf. [R, (4.1)]). Let X be a toric variety and D the complement of the big torus regarded as a reduced divisor. Then $K_X + D \sim 0$.

2.7 (the Kleiman-Mori Cone). Let $f : X \rightarrow Y$ be a proper morphism between normal varieties X and Y ; a 1-cycle of X/Y is a formal sum $\sum a_i C_i$ with complete curves C_i in the fibers of f , and $a_i \in \mathbb{Z}$. We put

$$Z_1(X/Y) := \{1\text{-cycles of } X/Y\},$$

and

$$Z_1(X/Y)_{\mathbb{Q}} := Z_1(X/Y) \otimes \mathbb{Q}.$$

There is a pairing

$$\text{Pic}(X) \times Z_1(X/Y)_{\mathbb{Q}} \rightarrow \mathbb{Q}$$

defined by $(\mathcal{L}, C) \mapsto \deg_C \mathcal{L}$, extended by bilinearity. Define

$$N^1(X/Y) := (\text{Pic}(X) \otimes \mathbb{Q}) / \equiv$$

and

$$N_1(X/Y) := Z_1(X/Y)_{\mathbb{Q}} / \equiv,$$

where the *numerical equivalence* \equiv is by definition the smallest equivalence relation which makes N^1 and N_1 into dual spaces.

Inside $N_1(X/Y)$ there is a distinguished cone of effective 1-cycles,

$$\text{NE}(X/Y) = \{ Z \mid Z \equiv \sum a_i C_i \text{ with } a_i \in \mathbb{Q}_{\geq 0} \} \subset N_1(X/Y).$$

A subcone $F \subset \text{NE}(X/Y)$ is said to be *extremal* if $u, v \in \text{NE}(X/Y)$, $u + v \in F$ imply $u, v \in F$. The cone F is also called an *extremal face* of $\text{NE}(X/Y)$. A one-dimensional extremal face is called an *extremal ray*.

We define the *relative Picard number* $\rho(X/Y)$ by

$$\rho(X/Y) := \dim_{\mathbb{Q}} N^1(X/Y) < \infty.$$

An element $D \in N^1(X/Y)$ is called *f-nef* if $D \geq 0$ on $\text{NE}(X/Y)$.

If X is complete and Y is a point, write $\text{NE}(X)$ and $\rho(X)$ for $\text{NE}(X/Y)$ and $\rho(X/Y)$, respectively. We note that $N_1(X/Y) \subset N_1(X)$, and $N^1(X/Y)$ is the corresponding quotient of $N^1(X)$.

2.2. Singularities of pairs. In this subsection, we quickly review the definitions of singularities which we use in the MMP. For details, see, for example, [KM, §2.3]. We recommend that the reader skip this subsection on first reading.

2.8. Let us recall the definitions of the singularities for pairs.

Definition 2.9 (Discrepancies and Singularities of Pairs). Let X be a normal variety and $D = \sum d_i D_i$ a \mathbb{Q} -divisor on X , where D_i are distinct and irreducible such that $K_X + D$ is \mathbb{Q} -Cartier. Let $f : Y \rightarrow X$ be a proper birational morphism from a normal variety Y . Then we can write

$$K_Y = f^*(K_X + D) + \sum a(E, X, D)E,$$

where the sum runs over all the distinct prime divisors $E \subset Y$, and $a(E, X, D) \in \mathbb{Q}$. This $a(E, X, D)$ is called the *discrepancy* of E with respect to (X, D) . We define

$$\text{discrep}(X, D) := \inf_E \{a(E, X, D) \mid E \text{ is exceptional over } X\}.$$

From now on, we assume that $0 \leq d_i \leq 1$ for every i . We say that (X, D) is

$$\begin{cases} \text{terminal} \\ \text{canonical} \\ \text{klt} \\ \text{plt} \\ \text{lc} \end{cases} \quad \text{if } \text{discrep}(X, D) \begin{cases} > 0, \\ \geq 0, \\ > -1 \text{ and } \lfloor D \rfloor = 0, \\ > -1, \\ \geq -1. \end{cases}$$

Here klt is an abbreviation for *Kawamata log terminal*, plt for *purely log terminal*, and lc for *log-canonical*.

If there exists a log resolution $f : Y \rightarrow X$ of (X, D) , that is, Y is non-singular, the exceptional locus $\text{Exc}(f)$ is a divisor, and $\text{Exc}(f) \cup f^{-1}(\text{Supp} D)$ is a simple normal crossing divisor, such that $a(E_i, X, D) > -1$ for every exceptional divisor E_i on Y , then the pair (X, D) is said to be dlt, which is an abbreviation for *divisorial log terminal*.

For details on dlt, see [KM, Definition 2.37, Theorem 2.44]. The following results are well-known to the experts. See, for example, [Fj1, Lemma 5.2] or [Ma, Proposition 14-3-2].

Proposition 2.10. *Let X be a toric variety and D the complement of the big torus regarded as a reduced divisor. Then (X, D) is log-canonical. Let $D = \sum_i D_i$ be the irreducible decomposition of D . We assume that $K_X + \sum_i a_i D_i$ is \mathbb{Q} -Cartier, where $0 \leq a_i < 1$ (resp. $0 \leq a_i \leq 1$) for every i . Then $(X, \sum a_i D_i)$ is klt (resp. log-canonical).*

3. TORIC MORI THEORY

We will work over an algebraically closed field k of arbitrary characteristic throughout this section. Let us explain the Minimal Model Program for toric varieties.

3.1 (Minimal Model Program for Toric Varieties). We start with a projective toric morphism $f : X \rightarrow Y$, that is, f is induced by a map of lattices, where $X =: X_0$ is a \mathbb{Q} -factorial toric variety, and a \mathbb{Q} -divisor $D_0 := D$ on X . The aim is to set up a recursive procedure that creates intermediate $f_i : X_i \rightarrow Y$ and D_i on X_i . After finitely many steps, we obtain final objects $\tilde{f} : \tilde{X} \rightarrow Y$ and \tilde{D} on \tilde{X} . Assume that we have already constructed $f_i : X_i \rightarrow Y$ and D_i with the following properties:

- (i) X_i is \mathbb{Q} -factorial and f_i is projective,
- (ii) D_i is a \mathbb{Q} -divisor on X_i .

If D_i is f_i -nef, then we set $\tilde{X} := X_i$ and $\tilde{D} = D_i$. If D_i is not f_i -nef, then we can take an extremal ray R of $\text{NE}(X_i/Y)$ such that $R \cdot D_i < 0$ (see Theorem 4.1 below). Thus we have a contraction morphism $\varphi_R : X_i \rightarrow W_i$ over Y (see Theorem 4.5 below). If $\dim W_i < \dim X_i$ (in which case we call φ_R a *Fano contraction*), then we set $\tilde{X} := X_i$ and $\tilde{D} := D_i$ and stop the process. If φ_R is birational and contracts a divisor (we call this a *divisorial contraction*), then we put $X_{i+1} := W_i$, $D_{i+1} := \varphi_{R*} D_i$ and repeat this process. In the case where φ_R is small (we call this a *flipping contraction*), then there exists a log-flip $\psi : X_i \dashrightarrow X_i^+$ over Y . Here, a log-flip means an *elementary transformation* with respect to D_i (see Theorem 4.8 below). Note that ψ is an isomorphism in codimension one. We put $X_{i+1} := X_i^+$, $D_{i+1} := \psi_* D_i$ and repeat this process. By counting the relative Picard number $\rho(X_i/Y)$, divisorial contractions can occur finitely many times (see Theorem 4.5). By Theorem 4.9 below, every sequence of log-flips terminates after finitely many steps. So, this process always terminates and we obtain $\tilde{f} : \tilde{X} \rightarrow Y$ and \tilde{D} . We call this process the *(D-)Minimal Model Program over Y*, where D is a divisor used in the process. When we apply the Minimal Model Program (MMP, for short), we say that, for example, we *run the MMP over Y with respect to the divisor D*.

Remark 3.2 (Toric Mori theory vs the general MMP). The general (log) MMP is still conjectural in dimension ≥ 4 (see [KM, §3.7]). As we see in Section 4, the MMP for toric varieties is fully established and it works for *any* \mathbb{Q} -divisor D . A generalization of the MMP for non- \mathbb{Q} -factorial toric varieties is treated in [Fj2, Section 2].

Remark 3.3 (Original toric Mori theory). In the above MMP, if we assume that Y is projective, X has only terminal singularities, f is birational, and $D = K_X$ the canonical divisor of X , then we can recover the original toric Mori theory in [R, (0.2) Theorem].

Remark 3.4. In 3.1, it is sufficient to assume that D is an \mathbb{R} -divisor. We do not treat the \mathbb{R} -generalization here because this generalization is obvious for experts and we do not need \mathbb{R} -divisors in this paper. We leave the details to the reader.

Remark 3.5. In general, the assumption that X is \mathbb{Q} -factorial is not crucial in the toric Mori theory. By [Fj1, Corollary 5.9], there always exists a small projective \mathbb{Q} -factorialization. Namely, for any toric variety X , there exists a small projective toric morphism $\tilde{X} \rightarrow X$ such that \tilde{X} is \mathbb{Q} -factorial. We refer the reader to [Fj1, Section 5], which is a baby version of this paper. We note that we can remove the assumption that X is complete in [Fj1, Theorem 5.5, Proposition 5.7] by the results in Sections 3 and 4 of this paper.

Remark 3.6. Let X be a toric variety and D a \mathbb{Q} -divisor on X . The assumption that D is \mathbb{Q} -Cartier can be removed in some cases. In fact, by replacing X by its small projective \mathbb{Q} -factorialization we can assume that D is \mathbb{Q} -Cartier. See the proof of Corollary 5.8.

4. PROOF OF THE TORIC MORI THEORY

In this section, we give a simple and non-combinatorial proof of the toric Mori theory.

Theorem 4.1 (the Cone Theorem). *Let $f : X \rightarrow Y$ be a proper toric morphism. Then the cone*

$$\mathrm{NE}(X/Y) \subset N_1(X/Y)$$

is a polyhedral convex cone. Moreover, if f is projective, then the cone is strongly convex.

Proof. By taking the Stein factorization of f , we may assume that f is surjective with connected fibers. We consider $V(\sigma) \subset Y$ for some cone σ . Then $f^{-1}(V(\sigma))$ is a union of $V(\tau) \subset X$ for some cones τ since f is a proper toric morphism. We divide Y into a finite disjoint union of tori $Y = \coprod_i Y_i$. We put $V_i := f^{-1}(Y_i) \rightarrow Y_i$. Let $\coprod_j V_{ij}$ be the normalization of V_i . Then we can check that V_{ij} is a toric variety for every i, j by using the above fact, that is, $f^{-1}(V(\sigma))$ is a union of orbit closures, inductively on $\dim V(\sigma)$. We note that V_{ij} is dominant onto Y_i for every i, j since Y_i is a torus. So, we obtain a collection of proper

surjective toric morphisms with connected fibers: $\{V_{ij} \longrightarrow Y_i\}_{i,j}$. By changing the notation V_{ij} , we write $\{f_i : X_i \longrightarrow Y_i\}_i$ for $\{V_{ij} \longrightarrow Y_i\}_{i,j}$. We note that $i \neq i'$ does not imply $Y_i \neq Y_{i'}$ in this notation. Since Y_i is a torus, $X_i \simeq F_i \times Y_i$ for every i , where F_i is a complete toric variety (cf. [Fl, p.41, Excercise]).

Claim. We have the following commutative diagram:

$$\begin{array}{ccc} N_1(F_i) & \simeq & N_1(X_i/Y_i) \\ \cup & & \cup \\ NE(F_i) & \simeq & NE(X_i/Y_i) \end{array}$$

for every i . In particular, $NE(X_i/Y_i)$ is a polyhedral convex cone for every i .

Proof of Claim. We consider the cycle map $Z_1(F_i) \longrightarrow Z_1(X_i/Y_i)$ which is induced by the inclusion $F_i \simeq F_i \times \{\text{a point of } Y_i\} \subset F_i \times Y_i \simeq X_i$. It induces

$$\varphi_i : N_1(F_i) \longrightarrow N_1(X_i/Y_i).$$

Let $0 \neq v \in N_1(F_i)$. Then there exists $\mathcal{L} \in \text{Pic}(F_i)$ such that $\mathcal{L} \cdot v \neq 0$. Let $p_i : X_i \longrightarrow F_i$ be the first projection. Then $p_i^* \mathcal{L} \cdot \varphi_i(v) = \mathcal{L} \cdot v \neq 0$ by the projection formula. Therefore, φ_i is injective. Since Y_i is a torus and $X_i \simeq F_i \times Y_i$, it is obvious that φ_i is surjective. Since $NE(F_i)$ is well-known to be a polyhedral convex cone (cf. [Fl, p.96 Proposition]), the other parts are obvious. \square

We consider the following commutative diagram:

$$\begin{array}{ccc} \bigoplus_i N_1(X_i/Y_i) & \twoheadrightarrow & N_1(X/Y) \\ \cup & & \cup \\ \bigoplus_i NE(X_i/Y_i) & \twoheadrightarrow & NE(X/Y). \end{array}$$

We note that $\bigoplus_i Z_1(X_i/Y_i) \longrightarrow Z_1(X/Y)$ is surjective. So, by combining it with the previous claim, we obtain the required cone theorem for $NE(X/Y) \subset N_1(X/Y)$. The last part follows from Kleiman's criterion. \square

We give another proof of Theorem 4.1, which works under the assumption that the characteristic of k is zero. This assumption is required only in the following proof.

Proof of Theorem 4.1 in characteristic zero. Assume that the characteristic of k is zero. First, we further assume that X is quasi-projective and \mathbb{Q} -factorial. Let T be the big torus of X . We put $D = \sum_i D_i =$

$X \setminus T$ regarded as a reduced divisor. We can take an ample \mathbb{Q} -divisor $L = \sum_i a_i D_i$ with $0 < a_i < 1$ for every i . Then

$$-(K_X + \sum_i (1 - a_i) D_i) \sim \sum_i a_i D_i$$

is ample (obviously, f -ample) and $(X, \sum_i (1 - a_i) D_i)$ is klt by Proposition 2.10. So, the well-known relative cone theorem (see, for example, [KMM, Theorem 4-2-1] or [KM, Theorem 3.25]) implies that $\text{NE}(X/Y)$ is a rational polyhedral convex cone. Next, by Chow's lemma and the desingularization theorem, we can take a proper birational toric morphism $X' \rightarrow X$ from a non-singular quasi-projective toric variety X' . So, the general case follows from the above special case. Details are left to the reader. \square

Remark 4.2. In Theorem 4.1, if X is complete, then every extremal ray of $\text{NE}(X/Y)$ is spanned by torus invariant curves on X . Related topics are treated in the first author's paper [Fj1]. If X is not complete, then $\text{NE}(X/Y)$ is not necessarily spanned by a torus invariant curve, as the following example shows:

Example 4.3. Let Y be a one-dimensional (not necessarily complete) toric variety. We put $X = Y \times \mathbb{P}^1$. Let $f : X \rightarrow Y$ be the first projection. Then $\text{NE}(X/Y)$ is a half line. When Y is a one-dimensional torus, there are no torus invariant curves in the fibers of f . If $Y \simeq \mathbb{P}^1$ or \mathbb{A}^1 , then $\text{NE}(X/Y)$ is spanned by a torus invariant curve in a fiber of f .

The following remark is obvious since a torus is a connected linear algebraic group (cf. [Su, Lemma 5]). We present it for the reader's convenience.

Remark 4.4. Let T be the big torus of X . Then T acts on $\text{NE}(X/Y)$. Let R be an extremal ray of $\text{NE}(X/Y)$. Then there exists a nef torus invariant Cartier divisor D on X such that $D \cdot [C] = 0$ if and only if $[C] \in R$. So, R is T -invariant. Thus, T acts on $\text{NE}(X/Y)$ trivially. Hence, the action of T on $N_1(X/Y)$ is trivial as well.

Therefore, an extremal ray R of $\text{NE}(X/Y)$ does not necessarily contain a torus invariant curve but is torus invariant.

Theorem 4.5 (the Contraction Theorem). *Let $f : X \rightarrow Y$ be a projective toric morphism. Let F be an extremal face of $\text{NE}(X/Y)$. Then there exists a projective surjective toric morphism*

$$\varphi_F : X \rightarrow Z$$

over Y with the following properties:

- (i) Z is a toric variety that is projective over Y .
- (ii) φ_F has connected fibers.
- (iii) Let C be a curve in a fiber of f . Then $[C] \in F$ if and only if $\varphi_F(C)$ is a point.

Furthermore, if F is an extremal ray R and X is \mathbb{Q} -factorial, then Z is \mathbb{Q} -factorial and $\rho(Z/Y) = \rho(X/Y) - 1$ if φ_R is not small.

Proof. Since $\text{NE}(X/Y)$ is a polyhedral convex cone, we can take an f -nef Cartier divisor D such that $D \cdot [C] \geq 0$ for every $[C] \in \text{NE}(X/Y)$ and $D \cdot [C] = 0$ if and only if $[C] \in F$. By replacing D by a linearly equivalent divisor, we may assume that D is a torus invariant Cartier divisor on X . We put $\varphi_F : X \rightarrow Z$ as in the the following Proposition 4.6. Then $\varphi_F : X \rightarrow Z$ has the required properties. The latter part is well-known. See, for example, [KM, Proposition 3.36]. \square

Proposition 4.6. *Let $f : X \rightarrow Y$ be a proper surjective toric morphism between toric varieties. Let D be an f -nef torus invariant Cartier divisor on X . Then D is f -free, that is, $f^*f_*\mathcal{O}_X(D) \rightarrow \mathcal{O}_X(D)$ is surjective. Moreover, we have a projective toric morphism $\varphi : X \rightarrow Z$ over Y such that*

- (i) φ has only connected fibers, and
- (ii) For any irreducible curve on X with $f(C)$ being a point, $\varphi(C)$ is a point if and only if $D \cdot C = 0$.

Proof. Let

$$f : X \xrightarrow{g} \tilde{Y} \xrightarrow{h} Y$$

be the Stein factorization of f . For the first part, we may assume that Y is affine (hence so is \tilde{Y}). It is sufficient to prove that D is g -free. We can apply the argument in [Fl, p.68, Proposition] with minor modifications. See also [N, Chapter IV. 1.8. Lemma (2)] and [Ma, Lemma 14-1-11]. For the second part, $\varphi : X \rightarrow Z := \text{Proj}_{\tilde{Y}} \bigoplus_{m \geq 0} g_*\mathcal{O}_X(mD)$ is equivariant by construction. When \tilde{Y} is a point, it is well-known that Z is a projective toric variety that is constructed from a suitable polytope. Let $T \subset \tilde{Y}$ be the big torus. Then $g^{-1}(T) \simeq T \times F$ for some complete toric variety F . So, Z contains a torus as a non-empty Zariski open set by the above case where \tilde{Y} is a point. It is obvious that Z is normal and has a suitable torus action by construction. Therefore, Z is the required toric variety. \square

Theorem 4.7 (Finitely Generatedness of Divisorial Algebra). *Let $f : X \rightarrow Y$ be a proper birational toric morphism and D a torus invariant*

Cartier divisor on X . Then

$$\bigoplus_{m \geq 0} f_* \mathcal{O}_X(mD)$$

is a finitely generated \mathcal{O}_Y -algebra.

Proof. We may assume that Y is affine. So, it is sufficient to show that

$$\bigoplus_{m \geq 0} H^0(X, \mathcal{O}_X(mD))$$

is a finitely generated k -algebra. We put $X = X(\Delta)$, that is, X is a toric variety associated to a fan Δ in N . Let ψ_D be the support function of D . We put

$$P_D = \{u \in M_{\mathbb{R}} \mid u \geq \psi_D \text{ on } |\Delta|\},$$

and

$$P_{aD} = \{u \in M_{\mathbb{R}} \mid u \geq a\psi_D \text{ on } |\Delta|\},$$

where $M := \text{Hom}_{\mathbb{Z}}(N, \mathbb{Z})$ is the dual lattice of N and $|\Delta|$ stands for the *support* of the fan Δ . We define

$$C = \{(u, a) \in M_{\mathbb{R}} \times \mathbb{R}_{\geq 0} \mid u \in P_{aD}\},$$

and $C_{\mathbb{Z}} = C \cap (M \times \mathbb{Z}_{\geq 0})$. The k -algebra $\bigoplus_{m \geq 0} H^0(X, \mathcal{O}_X(mD))$ is the semi-group ring associated to $C_{\mathbb{Z}}$. We can easily check that C is a finite intersection of half spaces, which are defined over the rational numbers, in $M_{\mathbb{R}} \times \mathbb{R}$. Therefore, C is a rational polyhedral convex cone. Thus, $C_{\mathbb{Z}}$ is a finitely generated semi-group. This implies that $\bigoplus_{m \geq 0} H^0(X, \mathcal{O}_X(mD))$ is a finitely generated k -algebra. \square

We will generalize the above theorem in Corollary 5.8 below.

Theorem 4.8 (Elementary Transformation). *Let $\varphi : X \rightarrow W$ be a small toric morphism and D a torus invariant \mathbb{Q} -Cartier divisor on X such that $-D$ is φ -ample. Let l be a positive integer such that lD is Cartier. Then there exists a small projective toric morphism*

$$\varphi^+ : X^+ := \text{Proj}_W \bigoplus_{m \geq 0} \varphi_* \mathcal{O}_X(mlD) \rightarrow W$$

such that D^+ is a φ^+ -ample \mathbb{Q} -Cartier divisor, where D^+ is the proper transform of D on X^+ . The commutative diagram

$$\begin{array}{ccc} X & \dashrightarrow & X^+ \\ & \searrow & \swarrow \\ & W & \end{array}$$

is called the elementary transformation (with respect to D).

Moreover, if X is \mathbb{Q} -factorial and $\rho(X/W) = 1$, then so is X^+ and $\rho(X^+/W) = 1$.

Proof. The first part is obvious by the above theorem and the construction of $\varphi^+ : X \rightarrow W$. See, for example, [K⁺, 4.2 Proposition] or [KM, Lemma 6.2]. The latter part is well-known. See, for example, [KM, Proposition 3.37]. \square

Theorem 4.9 (Termination of Elementary Transformations). *Let*

$$\begin{array}{ccccccc} X_0 & \dashrightarrow & X_1 & \dashrightarrow & X_2 & \dashrightarrow & \cdots \\ & \searrow & \swarrow \searrow & & \swarrow & & \\ & & W_0 & & W_1 & & \end{array}$$

be a sequence of elementary transformations of toric varieties with respect to a fixed \mathbb{Q} -Cartier divisor D . More precisely, the commutative diagram

$$\begin{array}{ccc} X_i & \dashrightarrow & X_{i+1} \\ & \searrow & \swarrow \\ & & W_i \end{array}$$

is the elementary transformation with respect to D_i , where $D_0 = D$ and D_i is the proper transform of D on X_i , for every i (see Theorem 4.8). Then the sequence terminates after finitely many steps.

Proof. Suppose that there exists an infinite sequence of elementary transformations. Let Δ be the fan corresponding to X_0 . Since the elementary transformations do not change one-dimensional cones of Δ , there exist numbers $k < l$ such that the composition $X_k \dashrightarrow X_{k+1} \dashrightarrow \cdots \dashrightarrow X_l$ is an identity. This contradicts the negativity in Lemma 4.10 below. \square

The following result is easy but very important. The proof is well-known. See, for example, [KM, Lemma 3.38].

Lemma 4.10 (the Negativity Lemma). *We consider a commutative diagram*

$$\begin{array}{ccc} & Z & \\ \swarrow & & \searrow \\ U & \dashrightarrow & V \\ \searrow & & \swarrow \\ & W & \end{array}$$

and \mathbb{Q} -Cartier divisors D and D' on U and V , respectively, where

- (1) $f : U \rightarrow W$ and $g : V \rightarrow W$ are birational morphisms between varieties,
- (2) $f_*D = g_*D'$,

- (3) $-D$ is f -ample and D' is g -ample,
- (4) $\mu : Z \longrightarrow U$, $\nu : Z \longrightarrow V$ are common resolutions.

Then $\mu^*D = \nu^*D' + E$, where E is an effective \mathbb{Q} -divisor and is exceptional over W . Moreover, if f or g is non-trivial, then $E \neq 0$.

5. ON THE ZARISKI DECOMPOSITION

In this section, we treat the Zariski decomposition on toric varieties. Since the MMP works for any divisors, it is obvious that the Zariski decomposition holds with no extra assumptions.

There are many variants of the Zariski decomposition. We adopt the following definition (cf. [KMM, Definition 7-3-5]) here.

Definition 5.1 (the Zariski Decomposition). Let $f : X \longrightarrow Y$ be a proper surjective morphism of normal varieties. An expression $D = P + N$ with \mathbb{R} -Cartier divisors D , P and N on X is called the *Zariski decomposition of D relative to f in the sense of Cutkosky-Kawamata-Moriwaki* (we write C-K-M for short) if the following conditions are satisfied:

- (1) P is f -nef,
- (2) N is effective, and
- (3) the natural homomorphisms $f_*\mathcal{O}_X(\lfloor mP \rfloor) \longrightarrow f_*\mathcal{O}_X(\lfloor mD \rfloor)$ are bijective for all $m \in \mathbb{N}$.

The divisors P and N are said to be the *positive* and *negative part* of D , respectively.

Definition 5.2 (Pseudo-effective Divisors). Let $f : X \longrightarrow Y$ be a projective morphism between varieties and D a \mathbb{Q} -Cartier divisor on X . Then D is *f -pseudo-effective* if there is an f -big (see [KM, Definition 3.22]) Cartier divisor A on X such that $nD + A$ is f -big for every $n \geq 0$.

Remark 5.3. Let $f : X \longrightarrow Y$ be a projective morphism between varieties and D a \mathbb{Q} -Cartier divisor on X . It is not difficult to see that if D is f -pseudo-effective then $nD + A$ is f -big for every $n \geq 0$ and any f -big Cartier divisor A (cf. [Mo, (11.3)]). In particular, if D is an effective divisor on X , then D is f -pseudo-effective. More generally, if there exists $m > 0$ such that $f_*\mathcal{O}_X(mD) \neq 0$, then D is f -pseudo-effective.

The following theorem is a slight generalization of [K, Proposition 5]. Related topics are in [N, Chapter IV §1]. Both [K] and [N] showed how to subdivide a given fan.

Theorem 5.4 (cf. [K, Proposition 5]). *Let $f : X \longrightarrow Y$ be a projective surjective toric morphism and D a \mathbb{Q} -Cartier divisor on X . Assume*

that D is f -pseudo-effective. Then there exists a projective birational toric morphism $\mu : Z \rightarrow X$ such that μ^*D has a Zariski decomposition relative to $f \circ \mu$ in the sense of C-K-M whose positive part is $f \circ \mu$ -semi-ample (see [KMM, Definition 0-1-4]).

Remark 5.5. If $f_*\mathcal{O}_X(mD) \neq 0$ for some positive integer m , then it is easy to check that $(f_k)_*\mathcal{O}_{X_k}(mD_k) \neq 0$ for every k , where $f_k : X_k \rightarrow Y$ is as in the following proof. Therefore, the following proof works without any changes even if we replace the assumption that D is f -pseudo-effective with a slightly stronger one that $f_*\mathcal{O}_X(mD) \neq 0$ for some positive integer m . Thus, we may not have to introduce the notion of f -pseudo-effective divisors. See Corollary 5.6 and the proof of Corollary 5.8 below.

Proof of Theorem 5.4. By taking a resolution of singularities, we may assume that X is non-singular without loss of generality. Run the MMP on X over Y with respect to D . We obtain a sequence of divisorial contractions and elementary transformations over Y :

$$X =: X_0 \dashrightarrow X_1 \dashrightarrow X_2 \dashrightarrow \cdots \dashrightarrow X_k \dashrightarrow X_{k+1} \dashrightarrow \cdots$$

Since D_k is pseudo-effective over Y for every k , there exists l such that D_l is nef over Y (for the definition of D_k , see 3.1). The reader may verify that relative pseudo-effectivity of D is preserved in each step by Lemma refneg and Remark 5.3. Take a non-singular quasi-projective toric variety Z with proper birational toric morphisms $\mu : Z \rightarrow X$ and $\mu_i : Z \rightarrow X_i$ for every $0 < i \leq l$. Then we obtain that $\mu^*D = \mu_l^*D_l + E$, where E is an effective \mathbb{Q} -divisor by the negativity in Lemma 4.10. This decomposition is the Zariski decomposition of D in the sense of C-K-M. \square

The next corollary is obvious by Theorem 5.4.

Corollary 5.6. *Let $f : X \rightarrow Y$ be a projective surjective toric morphism and D a \mathbb{Q} -Cartier divisor on X . Then D is f -pseudo-effective if and only if $f_*\mathcal{O}_X(mD) \neq 0$ for some positive integer m .*

Remark 5.7. There exist various generalizations of Theorem 5.4. We do not pursue such generalizations here. For example, the above theorem holds for a (not necessarily \mathbb{R} -Cartier) \mathbb{R} -divisor D , with suitable modifications. We leave the details to the reader.

The following result is a generalization of Theorem 4.7.

Corollary 5.8 (Finitely Generatedness of Divisorial Algebra II). *Let $f : X \rightarrow Y$ be a proper surjective toric morphism and D a (not*

necessarily \mathbb{Q} -Cartier) Weil divisor on X . Then

$$\bigoplus_{m \geq 0} f_* \mathcal{O}_X(mD)$$

is a finitely generated \mathcal{O}_Y -algebra.

Proof. By Remarks 3.5 and 3.6, we may assume that X is \mathbb{Q} -factorial. Hence, D is \mathbb{Q} -Cartier. By replacing X birationally, we may assume that f is projective. If $f_* \mathcal{O}_X(mD) = 0$ for every $m > 0$, then the claim is obvious. Therefore, we may assume that $f_* \mathcal{O}_X(mD) \neq 0$ for some $m > 0$, that is, D is f -pseudo-effective. Since by Theorem 5.4, there exists a projective birational toric morphism $\mu : Z \rightarrow X$ such that $\mu^* D$ has a Zariski decomposition with $f \circ \mu$ -semi-ample positive part, $\bigoplus_{m \geq 0} f_* \mathcal{O}_X(mD)$ is finitely generated. \square

6. APPLICATION TO HYPERSURFACE SINGULARITIES

In this section, we apply the toric Mori theory to the study of singularities. We will recover Ishii's results [I1]. We will work over the complex number field \mathbb{C} , throughout this section.

Let us recall the notion of *non-degenerate* hypersurface singularities quickly. For the details, see [I1].

Definition 6.1 (Non-degenerate Polynomials). For a polynomial $f = \sum_{\mathbf{m}} a_{\mathbf{m}} x^{\mathbf{m}} \in \mathbb{C}[x_0, x_1, \dots, x_n]$, where $x^{\mathbf{m}} = x_0^{m_0} x_1^{m_1} \cdots x_n^{m_n}$ for $\mathbf{m} = (m_0, m_1, \dots, m_n) \in \mathbb{Z}_{\geq 0}^{n+1}$, and a face γ of the Newton polytope $\Gamma_+(f)$ of f , denote $\sum_{\mathbf{m} \in \gamma} a_{\mathbf{m}} x^{\mathbf{m}}$ by f_{γ} . A polynomial f is said to be *non-degenerate* if for every compact face γ of $\Gamma_+(f)$ the $\partial f_{\gamma} / \partial x_i$ ($i = 0, \dots, n$) have no common zero on $(\mathbb{C}^{\times})^{n+1}$.

The following definitions are due to Ishii. See the introduction of [I1]. For the definitions of the singularities, see Definition 2.9.

Definition 6.2 (Minimal and Canonical Models). Let $(x \in X)$ be a germ of normal singularity on an algebraic variety. We call a morphism $\varphi : Y \rightarrow X$ a *minimal* (resp. *the canonical*) *model* of $(x \in X)$, if

- (1) φ is proper, birational,
- (2) Y has at most terminal (resp. canonical) singularities, and
- (3) K_Y is φ -nef (resp. φ -ample).

It is obvious that if a canonical model exists, then it is unique up to isomorphisms over X .

The next theorem is [I1, Theorem 2.3].

Theorem 6.3. *Let $X \subset \mathbb{C}^{n+1}$ be a normal hypersurface defined by a non-degenerate polynomial f . Then $(0 \in X)$ has both a minimal model and canonical model.*

Proof. Take a projective birational toric morphism $g : V \longrightarrow \mathbb{C}^{n+1}$ such that V is a non-singular toric variety and the proper transform X' of X on V is non-singular (see, for example, [I1, Proposition 2.2]). Run the MMP over \mathbb{C}^{n+1} with respect to $K_V + X'$. Then we obtain $\tilde{\varphi} : (\tilde{V}, \tilde{X}) \longrightarrow \mathbb{C}^{n+1}$ such that $K_{\tilde{V}} + \tilde{X}$ is $\tilde{\varphi}$ -nef. We note that the pair (\tilde{V}, \tilde{X}) has canonical singularities and \tilde{V} has at most terminal singularities. So \tilde{V} is non-singular in codimension two. Thus we obtain $K_{\tilde{X}} = (K_{\tilde{V}} + \tilde{X})|_{\tilde{X}}$. Therefore, $K_{\tilde{X}}$ is nef over X . It is not difficult to check that \tilde{X} has at most terminal singularities. Hence, this \tilde{X} is a minimal model of $(0 \in X)$ (see Definition 6.2). By using the relative base point free theorem (see, for example, [KMM, Theorem 3-1-1 and Remark 3-1-2(1)], or [KM, Theorem 3.24]), we obtain the canonical model of $(0 \in X)$. \square

Definition 6.4 (Dlt and Log-canonical Models). Let $(x \in X)$ be as in Definition 6.2. We call a morphism $\varphi : Y \longrightarrow X$ a *dlt* (resp. the *log-canonical*) *model* of $(x \in X)$, if

- (1) φ is proper birational,
- (2) (Y, E) is dlt (resp. log-canonical), where E is the reduced exceptional divisor of φ , and
- (3) $K_Y + E$ is φ -nef (resp. φ -ample).

It is obvious that if a log-canonical model exists, then it is unique up to isomorphisms over X . The notion of dlt models is new.

The next result is a slight generalization of [I1, Theorem 3.1]. The arguments in the following proof are more or less known to the experts of the MMP.

Theorem 6.5. *Let $X \subset \mathbb{C}^{n+1}$ be a normal hypersurface defined by a non-degenerate polynomial f . Then $(0 \in X)$ has both a minimal model and log-canonical model.*

Proof. Take a projective birational toric morphism $f_0 : V_0 \longrightarrow \mathbb{C}^{n+1}$ such that V_0 is a non-singular toric variety and the proper transform X_0 of X on V_0 is non-singular. We may assume that the reduced exceptional divisor E_0 of f_0 intersects X_0 transversally, that is, $E_0 \cup X_0$ is a simple normal crossing divisor on V_0 (see, for example, [I1, Proposition 2.2]). We note that f_0 is an isomorphism outside E_0 . Run the MMP over \mathbb{C}^{n+1} with respect to $K_{V_0} + X_0 + E_0$. Then we obtain a

sequence of divisorial contractions and elementary transformations

$$V_0 \dashrightarrow V_1 \dashrightarrow V_2 \dashrightarrow \cdots \dashrightarrow V_k \dashrightarrow V_{k+1} \dashrightarrow \cdots,$$

and the final object $\tilde{f} : \tilde{V} \longrightarrow \mathbb{C}^{n+1}$ has the property that $K_{\tilde{V}} + \tilde{X} + \tilde{E}$ is \tilde{f} -nef, where \tilde{X} is the proper transform of X_0 on \tilde{V} and \tilde{E} is the reduced \tilde{f} -exceptional divisor. We note that the exceptional locus of $V_i \longrightarrow \mathbb{C}^{n+1}$ is of pure codimension one for every i since \mathbb{C}^{n+1} is non-singular. So, \tilde{f} is an isomorphism outside \tilde{E} . Since $(\tilde{V}, \tilde{X} + \tilde{E})$ is dlt (cf. [KM, Corollary 3.44]), \tilde{V} is non-singular in codimension two around $\tilde{X} \cap \tilde{E}$ (cf. [KM, Corollary 5.55]). Thus, we obtain that $K_{\tilde{X}} + \tilde{E}|_{\tilde{X}} = (K_{\tilde{V}} + \tilde{X} + \tilde{E})|_{\tilde{X}}$ and $(\tilde{X}, \tilde{E}|_{\tilde{X}})$ is dlt (cf. [KM, Proposition 5.59]). We have to check that $\tilde{E}|_{\tilde{X}}$ is a reduced $\tilde{f}|_{\tilde{X}}$ -exceptional divisor. Let $\tilde{E} = \sum_i \tilde{E}_i$ be the irreducible decomposition. It is sufficient to show that $\tilde{f}(\tilde{E}_i) \subset \text{Sing}(X)$, where $\text{Sing}(X)$ is the singular locus of X . We write $K_{\tilde{V}} + \tilde{X} + \sum a_i \tilde{E}_i = \tilde{f}^*(K_{\mathbb{C}^{n+1}} + X)$. So, $\sum (1 - a_i) \tilde{E}_i$ is \tilde{f} -nef. If $\tilde{f}(\tilde{E}_i) \not\subset \text{Sing}(X)$, then $a_i < 1$. We note that (\mathbb{C}^{n+1}, X) is plt outside $\text{Sing}(X)$ (for the definition of *plt*, see Definition 2.9). This implies that $\tilde{f}(\tilde{E}_i) \subset \text{Sing}(X)$ for every i by [KM, Lemma 3.39]. Therefore, $(\tilde{X}, \tilde{E}|_{\tilde{X}})$ is a dlt model of $(0 \in X)$. By construction, $\tilde{f}|_{\tilde{X}}$ is an isomorphism outside $\tilde{E}|_{\tilde{X}}$. Since $K_{\tilde{V}} + \tilde{X} + \tilde{E}$ is nef over \mathbb{C}^{n+1} , we obtain a contraction morphism $\tilde{V} \longrightarrow V'$ over \mathbb{C}^{n+1} with respect to the divisor $K_{\tilde{V}} + \tilde{X} + \tilde{E}$ (cf. Theorem 4.5). Let X^\heartsuit be the normalization of the proper transform of \tilde{X} on V' . We put $E^\heartsuit := \mu_*(\tilde{E}|_{\tilde{X}})$, where $\mu : \tilde{X} \longrightarrow X^\heartsuit$. Then it is not difficult to see that $K_{X^\heartsuit} + E^\heartsuit$ is ample over X , E^\heartsuit is the reduced exceptional divisor of $X^\heartsuit \longrightarrow X$, and $K_{\tilde{X}} + \tilde{E}|_{\tilde{X}} = \mu^*(K_{X^\heartsuit} + E^\heartsuit)$. So, $(X^\heartsuit, E^\heartsuit)$ is the required log-canonical model of $(0 \in X)$. \square

Remark 6.6 (On Ishii's constructions). Answering our questions, Ishii informed us that the definition of \overline{E} in Claim 3.8 in the proof of [I1, Theorem 3.1] is not correct. She told us that \overline{E} should be defined as $\nu^*(K_{T_N(\Sigma_2)} + X(\Sigma_2) + E) - K_{\overline{X}}$, and then all discussions go well in the proof of [I1, Theorem 3.1]. Though we did not check the proof according to her corrected definition of \overline{E} , we see that our models coincide with her models. From now on, we freely use the notation in [I1].

Let F be the complement of the big torus of $T_N(\Sigma)$. Then the pair $(T_N(\Sigma), X(\Sigma))$ (resp. $(T_N(\Sigma), X(\Sigma) + F)$) has only canonical (resp. log-canonical) singularities. Therefore, by the arguments in Claim 2.8 in

[I1], it is not difficult to see that $m_{\mathbf{q}} \geq 0$ (resp. $m_{\mathbf{q}} \geq -1$) for every $\mathbf{q} \in \tilde{\Sigma}[1] \setminus \Sigma_0[1]$ (resp. $\mathbf{q} \in \tilde{\Sigma}[1] \setminus \Sigma_2[1]$) without the assumption that $D_{\mathbf{q}} \cap X(\tilde{\Sigma}) \neq \emptyset$ in Claim 2.8 (resp. Claim 3.5) in [I1]. Thus, we see that $(T_N(\Sigma_0), X(\Sigma_0))$ (resp. $(T_N(\Sigma_2), X(\Sigma_2) + E)$) has only canonical (resp. log-canonical) singularities. Ishii told us that she did not know the notion of singularities of pairs when she wrote [I1]. Therefore, $T_N(\Sigma_0) \simeq \text{Proj}_{\mathbb{C}^{n+1}} \bigoplus_{m \geq 0} g_* \mathcal{O}_V(m(K_V + X'))$, where g , V , and X' are as in the proof of Theorem 6.3, and $T_N(\Sigma_2) \simeq \text{Proj}_{\mathbb{C}^{n+1}} \bigoplus_{m \geq 0} f_{0*} \mathcal{O}_X(m(K_{V_0} + X_0 + E_0))$, where V_0 , X_0 , E_0 , and f_0 are as in the proof of Theorem 6.5. So, it is not difficult to see that the models constructed in [I1] coincide with ours. Details are left to the reader.

REFERENCES

- [C] C. Casagrande, Contractible classes in toric varieties, *Math. Z.* **243** (2003), no. 1, 99–126.
- [D] V.I. Danilov, The geometry of toric varieties, (Russian) *Uspekhi Mat. Nauk* **33** (1978), no. 2(200), 85–134, 247. English translation: *Russian Math. Surveys* **33** (1978), no.2, 97–154.
- [Fj1] O. Fujino, Notes on toric varieties from Mori theoretic viewpoint, *Tohoku Math. J.* **55** (2003), no. 4, 551–564.
- [Fj2] O. Fujino, Equivariant completions of toric contraction morphisms, preprint, 2003.
- [Fl] W. Fulton, *Introduction to toric varieties*, *Annals of Mathematics Studies*, **131**, The William H. Roever Lectures in Geometry, Princeton University Press, Princeton, NJ, 1993.
- [I1] S. Ishii, Minimal, canonical and log-canonical models of hypersurface singularities, *Birational algebraic geometry (Baltimore, MD, 1996)*, 63–77, *Contemp. Math.*, **207**, Amer. Math. Soc., Providence, RI, 1997.
- [I2] S. Ishii, The minimal model theorem for divisors of toric varieties, *Tohoku Math. J.* **51** (1999), no. 2, 213–226.
- [K] Y. Kawamata, The Zariski decomposition of log-canonical divisors, *Algebraic geometry, Bowdoin, 1985*, 425–433, *Proc. Sympos. Pure Math.*, **46**, Part 1, Amer. Math. Soc., Providence, RI, 1987.
- [KMM] Y. Kawamata, K. Matsuda, and K. Matsuki, Introduction to the Minimal Model Problem, in *Algebraic Geometry, Sendai 1985*, *Advanced Studies in Pure Math.* **10**, (1987) Kinokuniya and North-Holland, 283–360.
- [KKMS] G. Kempf, F. Knudsen, D. Mumford, and B. Saint-Donat, *Toroidal Embeddings I*, *Springer Lecture Notes Vol.* **339**, 1973.
- [KM] J. Kollár, and S. Mori, *Birational geometry of algebraic varieties*, *Cambridge Tracts in Mathematics*, Vol. **134**, 1998.
- [K⁺] J. Kollár et al., *Flips and Abundance for algebraic threefolds*, *Astérisque* **211**, Soc. Math. de France, 1992.
- [L] R. Laterveer, Linear systems on toric varieties, *Tohoku Math. J.* **48** (1996), 451–458.

- [Ma] K. Matsuki, *Introduction to the Mori program*, Universitext, Springer-Verlag, New York, 2002.
- [Mo] S. Mori, Classification of higher-dimensional varieties, Algebraic geometry, Bowdoin, 1985 (Brunswick, Maine, 1985), 269–331, Proc. Sympos. Pure Math., 46, Part 1, Amer. Math. Soc., Providence, RI, 1987.
- [N] N. Nakayama, Zariski-decomposition and Abundance, preprint, 2002, a revised and expanded version of RIMS-1142.
- [O] T. Oda, *Convex bodies and algebraic geometry*, An introduction to the theory of toric varieties, Translated from the Japanese, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)] **15**, Springer-Verlag, Berlin, 1988.
- [OP] T. Oda and H. S. Park, Linear Gale transforms and Gelfand-Kapranov-Zelevinskij decompositions, Tohoku Math. J. **43** (1991), 357–399.
- [R] M. Reid, Decomposition of toric morphisms, Arithmetic and geometry, Vol.II, 395–418, Progr. Math., **36**, Birkhäuser Boston, MA, 1983.
- [Sh] V. V. Shokurov, Letters of a bi-rationalist V. Mld's and the termination of log flips, preprint, 2003.
- [Su] H. Sumihiro, Equivariant completion, J. Math. Kyoto Univ. **14** (1974), 1–28.
- [W] J. Wisniewski, Toric Mori theory and Fano Manifolds, *Geometry of Toric Varieties*, Séminaires et Congrès **6**, (2002), 249–272 (electronic).

GRADUATE SCHOOL OF MATHEMATICS, NAGOYA UNIVERSITY, CHIKUSA-KU
NAGOYA 464-8602 JAPAN

E-mail address: fujino@math.nagoya-u.ac.jp

DEPARTMENT OF MATHEMATICS, TOKYO INSTITUTE OF TECHNOLOGY, 2-12-1
OH-OKAYAMA, MEGURO-KU, TOKYO 152-8551, JAPAN

E-mail address: hirosato@math.titech.ac.jp